



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Management of anti-colton(a) alloimmunisation in pregnancy: a case report

Kimmich, Nina ; Brand, B ; Hustinx, H ; Komarek, A ; Zimmermann, R

DOI: <https://doi.org/10.1111/tme.12287>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-123581>

Journal Article

Accepted Version

Originally published at:

Kimmich, Nina; Brand, B; Hustinx, H; Komarek, A; Zimmermann, R (2016). Management of anti-colton(a) alloimmunisation in pregnancy: a case report. *Transfusion medicine*, 26(2):150-152.

DOI: <https://doi.org/10.1111/tme.12287>

1 Management of anti-colton^a alloimmunisation in pregnancy: a case report

3 Kimmich N, Brand B, Hustinx H, Komarek A, Zimmermann R

5 Nina Kimmich, MD, Division of Obstetrics, University Hospital of Zurich, Switzerland

6 Brigitte Brand, MD, Division of Haematology, University Hospital of Zurich,
7 Switzerland

8 Hein Hustinx, MD, Division of Immunohaematology, Interregional Blood Transfusion
9 SRC Ltd., Switzerland

10 Adriana Komarek, MD, Division of Immunohaematology, Blood Transfusion Service
11 SRK, Zurich, Switzerland

12 Roland Zimmermann, MD, Division of Obstetrics, University Hospital of Zurich,
13 Switzerland

15 Corresponding author:

16 Nina Kimmich, MD

17 Division of Obstetrics, University Hospital of Zurich

18 Frauenklinikstrasse 10, 8091 Zurich

19 Switzerland

20 Tel: 0041-44-255 1111

21 E-Mail: Nina.Kimmich@usz.ch

26 Running title: Anti-Co^a alloimmunisation in pregnancy

Diagnosis and management of anti-Co^a alloimmunisation during pregnancy is a rare and challenging condition. Only five cases have been described up to now, four in the 1970s and one in 2008, but diagnosis and management has changed in the meantime (Smith, Stratton et al. 1970, Simpson 1973, McIntyre, Finigan et al. 1976, Fuhrmann, Kloppenburg et al. 1979, Michalewska, Wielgos et al. 2008). We here present such a case, its management and favourable outcome.

Alloimmunisation of a Co^a negative pregnant woman, carrying a Co^a positive fetus, may cause haemolytic disease in the fetus and newborn (HDFN) (Smith, Stratton et al. 1970, McIntyre, Finigan et al. 1976, Michalewska, Wielgos et al. 2008). Today, monitoring is performed by laboratory testing of antibody titer, although it is unknown whether antibody titration is helpful (de Haas, Thurik et al. 2015). Of greater importance are doppler flow measurements of the peak systolic velocity of the middle cerebral artery (MCA-PSV) (Zimmerman, Carpenter et al. 2002, Michalewska, Wielgos et al. 2008, Moise and Argoti 2012, de Haas, Thurik et al. 2015). In cases of fetal anaemia, intrauterine blood transfusion (IUT) by cordocentesis is established, challenged by the allocation of compatible blood (de Haas, Thurik et al. 2015).

A 32-year-old woman (blood group 0 RhD-, RhC-, Rhc+, RhE-, Rhe+, K-, Co^a-) was admitted to our hospital at 20+3 gestational weeks (gw) with an alloimmunisation against blood group antigen Co^a with increasing antibody titer. Antibody identification was performed by indirect antiglobulin testing (IAT) and enzyme (papain) testing using an in-house panel and Coombs and neutral cards (IAT/ID and papain/ID) (BioRad, Cressier, Switzerland) (Table 1a). Titration was performed in IAT/ID using a Co^a heterozygous test cell. A Co^a antigen determination of the fetus' father, using IAT/ID and non-commercial anti-Co^a and anti-Co^b sera, showed that he was Co^a homozygous. Monitoring of the fetus was performed every one to two weeks by MCA-PSV

measurements (Table 1b). Between 28 and 32 gw the MCA-PSV gradually increased with values above the 95th percentile and a Monocyte Monolayer Assay (MMA) showed a value of 29% (Nance, Nelson et al. 1989), so that anaemia was suspected and fetal transfusion planned. We used Co^a-, RhD-, RhC-, RhE- and Kell negative blood from a donor, as women between 0 and 50 years of age should be transfused with Rh/K phenotype compatible blood only, according to the Swiss recommendations. We did not use blood from the mother, as autologous blood donation during pregnancy is not a common, but rarely performed procedure in Switzerland, especially in order to avoid lowering the mother's haemoglobin level. At 32+4 gw a cordocentesis was performed. An immediate intrauterine transfusion was initiated already before the fetal blood result was present, in order to avoid a possible second cordocentesis due to a dislocated needle. Knowing the haemoglobin/haematocrit of the fetus and of the preserved blood for transfusion, one can calculate the amount of blood needed to reach an appropriate fetal haemoglobin/haematocrit, according to the reference values by Mari (Mari, Deter et al. 2000). As the intraoperative blood results showed a haemoglobin and haematocrit of 123 g L⁻¹ and 35.5 %, respectively, transfusion was stopped. So far, 40 ml of RhD-, RhC-, RhE-, Kell- and Co^a negative, irradiated and washed red blood cells, depleted for plasma and leucocytes, with a haematocrit of 81.3 % were transfused, leading to a posttransfusion haemoglobin level and haematocrit of 168 g L⁻¹ and 48.7 %, respectively. MCA-PSV decreased to normal values within minutes and remained stable (Table 1b). Delivery was performed at 37+2 gw by elective cesarean as part of our standard protocol for alloimmunised pregnancies. Blood testing of the newborn confirmed blood group A RhD negative with antibodies against Co^a, detected by direct antiglobulin testing (DAT) (agglutination ++, eluate testing positive). Fetal umbilical cord haemoglobin and haematocrit levels were 156 g L⁻¹ and 47%, respectively (Table 1a). Due to fetal hyperbilirubinaemia on the fifth day postpartum, the neonate was

78 treated with phototherapy for three days (Table 1a). This was the reason, why mother
79 and child were discharged from hospital not before eight days postpartum, both without
80 complications.

81 Given the limited data on the relevance of antibody titer, close monitoring of the mother
82 and fetus was justified. Repeated antibody screening was performed for early detection
83 of a titer increase. Although a significant increase in antibody titer was documented,
84 this did not reflect fetal anaemia. Unfortunately, a clear cutoff for anti- Co^a antibody
85 titers, leading to fetal anaemia, is lacking (de Haas, Thurik et al. 2015). In our and other
86 cases of severe fetal anaemia, the antibody titer was measured at a level of 1:128 in
87 IAT and in a case of mild HDFN the titer was only 1:32 (Simpson 1973, McIntyre,
88 Finigan et al. 1976, Michalewska, Wielgos et al. 2008). Here, however, it is important
89 to consider the different methods of antibody titer determination used in the 1970s and
90 nowadays. Additionally, a MMA can be performed to evaluate the clinical significance
91 of the antibody. Monocyte reactivity greater than 20% was initially reported to correlate
92 with needs for transfusion (Nance, Arndt et al. 1987). Subsequent studies from the
93 same author, however, indicated that the assay may not always predict the outcome
94 of the infant, as seen in our case, and should not be used to evaluate HDFN. Before
95 diagnosing fetal anaemia by MCA-PSV, bilirubin extinction in amniotic fluid from
96 amniocentesis was performed (Moise and Argoti 2012, de Haas, Thurik et al. 2015).
97 This method was used in the four cases in the 1970s, but required repeated
98 amniocentesis with the risks of an invasive procedure (Simpson 1973, McIntyre,
99 Finigan et al. 1976, Fuhrmann, Kloppenburg et al. 1979). Measurement of the MCA-
100 PSV is the gold standard nowadays and was performed in our and Michalewska`s case
101 (Mari, Adrignolo et al. 1995, Michalewska, Wielgos et al. 2008). Compared to that case,
102 where MoM-values for the MCA-PSV cutoff were used, we used the reference values

published by Kurmanavicius (Mari, Adrignolo et al. 1995, Kurmanavicius, Streicher et al. 2001). However, in our case MCA-PSV values indicative for fetal anaemia also exceeded the MoM-values by Mari. No neonatal exchange or top-up transfusion was required in our case and in the ones of McIntyre and Smith, but in the other cases (Table 2). Although MCA-PSV values were indicative for fetal anaemia in our case, blood values were almost normal for the age of gestation.

The neonatal hyperbilirubinaemia, treated with phototherapy for three days, was probably not due to alloimmunisation against Co^a, but in the context of intensified postpartum haemolysis.

A great challenge in anti-Co^a alloimmunised patients is the allocation of compatible blood for transfusion. In Switzerland, this can be achieved by finding a donor through the “DGTI Rare Donor Register” (<http://www.iblutspende.ch/rare-donors.html>), by autologous donation of maternal blood or by transfusion of washed maternal erythrocytes (not a common procedure, limited to rare cases with missing compatible blood donors). In this case, the possible selection was even smaller due to the 0 RhD negative blood group of the patient (only 6% of blood donors). Only seven donors were compatible for the constellation RhD-, RhC, RhE-, K- and Co^a negative in Switzerland and only three of them available at that time. Furthermore, preparation time of two to three days had to be taken into consideration. Unfortunately, non-invasive treatment options are not available up to date.

Anti-Co^a alloimmunisation during pregnancy is a challenging situation. Both national and international blood donor registries are extremely helpful to identify compatible blood donors. Despite increased antibody concentrations, MCA-PSV and MMA, this case did not develop fetal or neonatal anaemia and did not require exchange transfusion after birth. Serologic methods can help to initiate fetal monitoring and

doppler MCA-PSV measurements might be helpful to avoid serial amniocentesis or fetal blood samplings, but can also be misleading in some cases.

Tables

154 Table 1:

155 a) Haematological course of mother and fetus/neonate

gestational weeks	anti-Co ^a titer	maternal haemoglobin (g/100mL)	maternal haematocrit (%)	fetal/neonatal haemoglobin (g/100mL)	fetal/neonatal haematocrit (%)	Neonatal blood bilirubin (mcmmol/L)	neonatal transcutaneous bilirubin (mcmmol/L)
6+3	2						
10+2	8	121	36				
13+2	16						
16+3	32						
20+3	128	129	37.6				
30+3	16						
32+4 (before transfusion)		127	37.5	123	35.5		
32+4 (after transfusion)				168	48.7		
37+1		122	35.3				
37+2 (cesarean)				156	47	39	59
1st day postpartum		117	35.8		56	105	109
2nd day postpartum					56	200	179
3rd day postpartum							225
4th day postpartum					55	276	241
5th day postpartum					56	300	
6th day postpartum					55	281	
7th day postpartum					61	299	
8th day postpartum					57	258	

156

157 b) MCA-PSV values in the course of pregnancy

gestational weeks	MCA -PSV (cm/s)
20+3	27
22+3	32
24+3	39
26+3	45
28+3	54*
29+3	60*
29+6	60*
30+3	65*
30+5	61*
31+3	62*
31+6	74*
32+2	72*
32+4 (before transfusion)	72*
32+4 (after transfusion)	43
32+5	54
33+5	63
34+6	65
36+4	72

158 *Value above the 95th percentile of reference limits

159

160

161 Table 2: Course and outcome of the six cases

Case	Anit-Co ^a titer	Amniocentesis (n)	Cord blood sampling (n)	Fetal anaemia	Intrauterine transfusion (n)	Neonatal anaemia	neonatal transfusion (n)
Smith 1970		0	0	none	0	none	0
Simpson 1973	128	2	0	severe	0	severe	2
McIntyre 1976	32	1	0	none	0	none	0
Fuhrmann 1979		2	0	mild	0	severe*	2*
Michalewska 2008	128	0	3	severe	1	severe	5
Our case	128	0	1	mild	1	mild	0

*due to bleeding out of a ruptured of vessel in presence of insertio velamentosa

Acknowledgements

There are no acknowledgements.

Contribution of the authors:

Nina Kimmich: Management and treatment of the patient, wrote the manuscript

Brigit Brand: Management of the patient, correction of the manuscript

Hein Hustinx: Management of the patient, correction of the manuscript

Adriana Komarek: Management of the patient, correction and translation of the manuscript

Roland Zimmermann: Management and treatment of the patient, correction of the manuscript

Disclosure of Interests

The authors of this manuscript have no conflicts of interest to disclose as described by

Transfusion Medicine. No funding was obtained.

References

184 de Haas, M., F. F. Thurik, J. M. Koelewijn and C. E. van der Schoot (2015). "Haemolytic
 185 disease of the fetus and newborn." Vox Sang.

186 Fuhrmann, U., W. Kloppenburg and H. W. Kruger (1979). "[Delivery of a pregnant
 187 woman with a rare phenotype in the Colton-blood group system (author's transl)]."
 188 Geburtshilfe Frauenheilkd **39**(1): 66-67.

189 Kurmanavicius, J., A. Streicher, E. M. Wright, J. Wisser, R. Muller, P. Royston, R.
 190 Huch, A. Huch and R. Zimmermann (2001). "Reference values of fetal peak systolic
 191 blood flow velocity in the middle cerebral artery at 19-40 weeks of gestation."
 192 Ultrasound Obstet Gynecol **17**(1): 50-53.

193 Mari, G., A. Adrignolo, A. Z. Abuhamad, J. Pirhonen, D. C. Jones, A. Ludomirsky and
 194 J. A. Copel (1995). "Diagnosis of fetal anemia with Doppler ultrasound in the pregnancy
 195 complicated by maternal blood group immunization." Ultrasound Obstet Gynecol **5**(6):
 196 400-405.

197 Mari, G., R. L. Deter, R. L. Carpenter, F. Rahman, R. Zimmerman, K. J. Moise, Jr., K.
 198 F. Dorman, A. Ludomirsky, R. Gonzalez, R. Gomez, U. Oz, L. Detti, J. A. Copel, R.
 199 Bahado-Singh, S. Berry, J. Martinez-Poyer and S. C. Blackwell (2000). "Noninvasive
 200 diagnosis by Doppler ultrasonography of fetal anemia due to maternal red-cell
 201 alloimmunization. Collaborative Group for Doppler Assessment of the Blood Velocity
 202 in Anemic Fetuses." N Engl J Med **342**(1): 9-14.

203 McIntyre, C., L. Finigan and A. L. Larsen (1976). "Anti-Coa implicated in hemolytic
 204 disease of the newborn." Transfusion **16**(1): 76-78.

205 Michalewska, B., M. Wielgos, B. Zupanska and R. Bartkowiak (2008). "Anti-Co(a)
 206 implicated in severe haemolytic disease of the foetus and newborn." Transfus Med
 207 **18**(1): 71-73.

208 Moise, K. J., Jr. and P. S. Argoti (2012). "Management and prevention of red cell
 209 alloimmunization in pregnancy: a systematic review." Obstet Gynecol **120**(5): 1132-
 210 1139.

211 Nance, S. J., P. Arndt and G. Garratty (1987). "Predicting the clinical significance of
 212 red cell alloantibodies using a monocyte monolayer assay." Transfusion **27**(6): 449-
 213 452.

214 Nance, S. J., J. M. Nelson, J. Horenstein, P. A. Arndt, L. D. Platt and G. Garratty (1989).
 215 "Monocyte monolayer assay: an efficient noninvasive technique for predicting the
 216 severity of hemolytic disease of the newborn." Am J Clin Pathol **92**(1): 89-92.

217 Simpson, W. K. (1973). "Anti-Coa and severe haemolytic disease of the newborn." S
 218 Afr Med J **47**(29): 1302-1304.

219 Smith, D. S., F. Stratton, P. Howell and R. Riches (1970). "An example of anti-Coa
 220 found in pregnancy." Vox Sang **18**(1): 62-66.

221 Zimmerman, R., R. J. Carpenter, Jr., P. Durig and G. Mari (2002). "Longitudinal
 222 measurement of peak systolic velocity in the fetal middle cerebral artery for monitoring
 223 pregnancies complicated by red cell alloimmunisation: a prospective multicentre trial
 224 with intention-to-treat." BJOG **109**(7): 746-752.

225